



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\therefore p = \frac{a[(\sin\theta' - \cos\theta') + (\sin\theta'' - \cos\theta'') + (\sin\theta''' - \cos\theta''') + \text{etc.}]}{a[(\sin\theta' + \cos\theta') + (\sin\theta'' + \cos\theta'') + (\sin\theta''' + \cos\theta''') + \text{etc.}]}$$

$$= \frac{a \int (\sin\theta - \cos\theta) d\theta}{a \int (\sin\theta + \cos\theta) d\theta}.$$

The limits of θ for favorable cases are $\frac{1}{2}\pi$ and $\frac{3}{2}\pi$ and doubled, for the total cases $\frac{1}{2}\pi$ and 0.

$$\therefore p = \frac{2 \int_{\frac{1}{2}\pi}^{\frac{3}{2}\pi} [\sin\theta - \cos\theta] d\theta}{\int_0^{\frac{1}{2}\pi} [\sin\theta + \cos\theta] d\theta} = (\sqrt{2} - 1).$$

NOTE.—The answer given in Professor Byerly's Integral Calculus, Edition of 1890, is $\frac{1}{2} - 1/\pi \log 2$.

This is the answer to the following problem: *From a point taken at random in the side of a square, a line is drawn at random across the square. What is the chance that the line will intersect the opposite side of the square?*

Professor Zerr interpreted the problem in this way and solved it accordingly.

Solutions somewhat similar to the one above were received from J. M. Colaw and Lon C. Walker.

Professor Walker sent a solution of No. 93. His answer is $\frac{1}{2}\pi p^3$. This difference in results is due to the different interpretations of the problem by Professors Zerr and Walker.

97. Proposed by L. C. WALKER, Assistant Professor of Mathematics, Leland Stanford Jr. University, Palo Alto, Cal.

A straight line is drawn at random across a circle, and five points are taken at random in the surface of the circle. Required the chance that all the points are on the same side of the line.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let θ = angle subtended by random chord at the center. Then area of the smaller segment is $a^2(\theta - \sin\theta\cos\theta)$. Hence the required chance is

$$p = \frac{\int_0^\pi 2[a^2(\theta - \sin\theta\cos\theta)]^5 a \sin\theta d\theta}{\int_0^\pi (\pi a^2)^5 a \sin\theta d\theta}$$

$$= \frac{1}{\pi^5} \int_0^\pi (\theta - \sin\theta\cos\theta)^5 \sin\theta d\theta = 1 - \frac{128}{9\pi^2} + \frac{2768896}{33075\pi^4} = .4184.$$